

DRAFT

Fire Regime Condition Class (FRCC) Interagency Handbook Reference Conditions

Modeler: Mike Schafale

Date: June 3, 2004

PNVG Code: AWPS

Potential Natural Vegetation Group: Atlantic Wet Pine Savannas

Geographic Area: Southeastern Virginia to Georgia

Description: Wet woodlands/savannas on wet mineral soils. Canopy dominated by *Pinus palustris*, sometimes mixed with *Pinus serotina*. There is generally little or no understory in reference condition, but a variety of hardwoods may occur with infrequent fire. The ground cover is dense and generally diverse. Grasses such as *Aristida stricta*, *Sporobolus pinetorum*, *Ctenium aromaticum*, and *Sporobolus teretifolius* dominate, and a large number of other grasses, sedges, and forbs including insectivorous plants are present. Shrubs are of fairly low density in reference condition but can become dense with fire exclusion.

Canopy trees are patchy in distribution, with regeneration in canopy gaps of ¼ acre or less in size, mid-successional clumps in similar size patches, and the oldest trees occurring as isolated individuals. The reference condition classes are aggregates of numerous patches well dispersed over the landscape. Canopy gaps are created by fire mortality, lightning, and wind throw at the scale of individual trees or several trees. Because of the irregular seed production of longleaf pine, canopy gaps may lack regeneration for several years.

This PNVG is distinguished from other longleaf pine-dominated by the presence of wetland herbs and shrubs. It includes the wet pine flatwoods of the Carolinas but not the *Serenoa repens*-containing flatwoods of the Gulf Coast region. It is abundant on remaining natural lands in the outer and middle Coastal Plain and occurs in small patches in the Fall Line Sandhills region.

Uncharacteristic vegetation types include even-aged canopy stands in which age structure has been homogenized by logging or clearing, examples where loblolly or slash pine have replaced some or all of the longleaf pine, examples where shrubs have become dense due to inadequate burning, and examples where the grass-dominated ground cover has been lost due to soil disturbance or past canopy closure. Full restoration to reference condition may take a number of burns, and may take many years if older trees are not present, but fire produces substantial ecological benefits before full restoration.

Fire Regime Description: Frequent surface fires, every 2-5 years, generally burn across large expanses. Fires are usually low intensity overall but will occasionally kill young regeneration patches and rarely kill individual older trees. Replacement fires in this model represent small patch mortality occurring in the context of extensive low intensity fires, rather than extensive stand replacement. Mosaic fire in the model represents the probability of a series of surface fires sufficient to move closed vegetation to open; single fires are generally ineffective.

Vegetation Type and Structure

Class*	Percent of Landscape	Description
A: canopy gaps	16	Canopy gaps, most single tree to quarter acre size, with pine regeneration up to 15 years old or lacking pine regeneration

		because no mast year has occurred since the gap opened. Native grassy ground cover dominated by <i>Aristida stricta</i> . Tree cover 0 to 50%.
B: middle-aged canopy with shrubs	4	Patches, mostly ¼ acre or less, with canopy pines 15-75 years old, with a substantial component of mid-story hardwoods or of shrubs encroaching in the absence of fire. Hardwood/shrub cover greater than 50%. Canopy pine cover 25-75%.
C: middle-aged canopy with herbs	38	Patches, most ¼ acre or less, with canopy pines 15-75 years old, with little hardwood component and only sparse shrubs due to frequent fire. <i>Aristida stricta</i> -dominated ground cover. Canopy pine cover 25-75%.
D: old canopy with herbs	39	Patches, most ¼ acre or less, with canopy pines 75 or more years old, with little hardwood component and only sparse shrubs due to frequent fire. <i>Aristida stricta</i> -dominated ground cover. Canopy pine cover 25-75%.
E: old canopy with shrubs	3	Patches with canopy pines 75 or more years old, with a substantial component of hardwoods and/or shrubs in either the overstory or understory. Ground cover shrubby or sparse. Hardwood/shrub cover greater than 50%.
Total	100	

*Formal codes for classes A-E are: AESP, BMSC, CMSO, DLSO, and ELSC, respectively.

Fire Frequency and Severity

Fire Severity	Fire Frequency (yrs)	Probability	Percent, All Fires	Description
Replacement Fire	100	.01	3%	Most replacement is in class A. Older pines are very fire-resistant and mortality is uncommon.
Non-Replacement Fire	3	.29	97%	Low intensity surface fires in all classes.
All Fire Frequency*	3	.30	100	

*All Fire Probability = sum of replacement fire and non-replacement fire probabilities. All Fire Frequency = inverse of all fire probability (previous calculation).

Modeling Assumptions:

Primary dynamic is the gap phase regeneration of longleaf pine. The model classes are small patches widely interspersed on the landscape. Replacement means death of longleaf pines as single trees or small clumps.

Most replacement fires occur in the earliest stage (class A). Older trees are very resilient to fire. Secondary dynamic (closed vs. open path) is the invasion of shrubs and hardwood trees in patches that escape fire.

Once shrubs are established, they slightly decrease probability of fire, but increase the probability that fires will kill the canopy pines.

Once established, shrubs are not easily eliminated by single fires, but may sometimes be eliminated by multiple fires. We have simulated this by using mosaic fire to represent the last of a series of surface fires that eliminates invading hardwoods without killing canopy pines.

References

Boyer, W. D. 1993. Long-term development of regeneration under pine seed tree and shelter wood stands. *Southern Journal of Applied Forestry* 17: 10-15.

- Brewer, J. S., and Platt, W. J. 1994a. Effects of fire season and herbivory on reproductive success in a clonal forb, *Pityopsis graminifolia*. *Journal of Ecology* 82: 665-675.
- Brewer, J. S., and Platt, W. J. 1994b. Effects of fire season and soil fertility on clonal growth in the pyrophilic forb, *Pityopsis graminifolia* (Asteraceae). *American Journal of Botany* 81: 805-814.
- Brown, James K.; Smith, Jane Kapler, eds. 2000. Wildland fire in ecosystems: effects of fire on flora. Gen. Tech. Rep. RMRS-GTR-42-vol. 2. Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 257 p.
- Christensen, N. L. 1981. Fire regimes in southeastern ecosystems. Pages 112-136 in H. A. Mooney, T. M. Bonnicksen, N. L. Christensen, J. E. Lotan, and W. A. Reiners (eds.). Fire regimes and ecosystem properties. USDA Forest Service General Technical Report WO-26.
- Crocker, T. C., and Boyer, W. D. 1984. Regenerating longleaf pine naturally. USDA Forest Service Research Paper SE-105.
- Glitzenstein, J. S., Harcombe, P. A., and Streng, D. R. 1986. Disturbance, succession, and maintenance of species diversity in an east Texas forest. *Ecological Monographs* 56: 243-258.
- Hains, M. J., Mitchell, R. J., Palik, B. J., and Boring, L. R. 1999. Abundance and distribution of native legumes (*Leguminosae*) in frequently burned longleaf pine (*Pinaceae*)—wiregrass (*Poaceae*) ecosystems. *American Journal of Botany* 86:1606-1614.
- Hermann, S. M., Van Hook, T., Flowers, R. W., Brennan, L. A., Glitzenstein, J. S., Streng, D. R., Walker, J. L., Myers, R. L. 1998. Fire and biodiversity: studies of vegetation and arthropods. North American Wildland and Natural Resource Conference 63:384-401.
- Hiers, J. K., Wyatt, R., Mitchell, R. J. 2000. The effects of fire regime on legume reproduction in longleaf pine savannas: is a season selective? *Oecologia* 125:521-530.
- Jacqmain, E.I., Jones R.H, and Mitchell, R.M. 1999. Influences of frequent cool-season burning across a soil moisture gradient on oak community structure in longleaf pine ecosystems. *American Midland Naturalist* 141: 85-100.
- Kirkman, L. K., Drew, M. B., Edwards, D. 1998. Effects of experimental fire regimes on the population dynamics of *Schwalbea americana* L. *Plant Ecology* 137:115-137.
- Palik, B. J., and Pederson, N. 1996. Natural disturbance and overstory mortality in longleaf pine ecosystems. *Canadian Journal of Forest Research* 26: 2035-2047.
- Palik, B. J., Mitchell, R. J., Houseal, G., and Pederson, N. 1997. Effects of canopy structure on resource availability and seedling responses in a longleaf pine ecosystem. *Canadian Journal of Forest Research* 27: 1458-1464.
- Palik, B. J., and Engstrom, R. T. 1999. Species composition. Pages 65-94 in M. L. Hunter, Jr. (ed.). Maintaining biodiversity in forest ecosystems. Cambridge University Press.

- Peet, R. K., and Allard, D. J. 1993. Longleaf pine vegetation of the Southern Atlantic and Eastern Gulf Coast regions: a preliminary classification. In Proceedings of the 18th Tall Timbers Fire Ecology Conference. Tallahassee, Florida, USA.
- Platt, W. J., and Rathbun, S. L. 1993. Populations dynamics of an old-growth population of longleaf pine (*Pinus palustris*). In Proceedings of the 18th Tall Timbers Fire Ecology Conference. Tallahassee, Florida, USA.
- Platt, W. J., Evans, G. W., and Rathbun, S. L. 1988. The population dynamics of a long-lived conifer (*Pinus palustris*). *American Naturalist* 131: 491-525.
- Platt, W. J., Evans, G. W., Davis, M. M. 1988. Effects of fire season on flowering of forbs and shrubs in longleaf pine forests. *Oecologia* 76: 353-363.
- Schmidt, Kirsten M, Menakis, James P., Hardy, Colin C., Hann, Wendel J., Bunnell, David L. 2002. Development of coarse-scale spatial data for wildland fire and fuel management. Gen. Tech. Rep. RMRS-GTR-87. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 41 p. + CD.
- Streng, D, R, Glitzenstein, J.S., Platt, W. J. 1993) Evaluating effects of season of burn in longleaf pine forests: A critical literature review and some results from an ongoing long-term study. Proceedings of the Tall Timbers Fire Ecology Conference 18:227-259.
- Robbins, L. E., and Myers, R. L. 1992. Seasonal effects of prescribed burning in Florida. Miscellaneous Publications of the Tall Timbers Research Station 8: 1-97.
- U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (2002, December). Fire Effects Information System, [Online]. Available: <http://www.fs.fed.us/database/feis/>.
- Walker, J., and Peet, R. K. 1983. Composition and species diversity of pine-wiregrass savannas of the Green Swamp, North Carolina. *Vegetatio* 55: 163-179.
- Walters, J. R. 1991. Application of ecological principals to the management of endangered species: the case of the red-cockaded woodpecker. *Annual Review of Ecology and Systematics* 22: 505-523.
- Ware, S., Frost, C., and Doerr, P. D. 1993. Southern mixed hardwood forest: The former longleaf pine forest. Pages 447-493 in W.H. Martin, S.G. Boyce, and A.C. Echternacht (eds.). Biodiversity of the southeastern United States. John Wiley and Sons, Inc., New York, New York, USA.

PERSONAL COMMUNICATION (if applicable):

VDDT File Documentation

Include screen captures (print-screens) from any of the VDDT graphs that were used to develop reference conditions.









